

IN THE CLAIMS

This listing of the claims replaces all prior versions of the claims in the application.

1. (Withdrawn.) An isolated polypeptide selected from the group consisting of:

- a) a polypeptide comprising an amino acid sequence of SEQ ID NO:1,
- b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to an amino acid of SEQ ID NO:1,
- c) a biologically active fragment of a polypeptide having an amino acid sequence of SEQ ID NO:1, and
- d) an immunogenic fragment of a polypeptide having an amino acid sequence of SEQ ID NO:1.

2. (Withdrawn.) An isolated polypeptide of claim 1 having an amino acid sequence of SEQ ID NO:1.

3. (Currently Amended.) An isolated polynucleotide encoding a polypeptide selected from the group consisting of:

- a) a polypeptide comprising the amino acid sequence of SEQ ID NO:1,
- b) a polypeptide comprising a naturally occurring amino acid sequence at least 90% identical to the amino acid of SEQ ID NO:1,
- c) a biologically active fragment of a polypeptide having the amino acid sequence of SEQ ID NO:1, wherein said biologically active fragment has serine dehydratase hydratase activity, and
- d) an immunogenic fragment of a polypeptide having the amino acid sequence of SEQ ID NO:1, wherein said immunogenic fragment generates an antibody that specifically binds to SEQ ID NO:1.

4. (Previously presented.) An isolated polynucleotide of claim 3 encoding the amino acid sequence of SEQ ID NO:1.

5. (Previously presented.) An isolated polynucleotide of claim 4 having the polynucleotide sequence of SEQ ID NO:2.

6. (Original.) A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

7. (Original.) A cell transformed with a recombinant polynucleotide of claim 6.

8. (Withdrawn.) A transgenic organism comprising a recombinant polynucleotide of claim 6.

9. (Currently Amended.) A method of producing a polypeptide selected from the group consisting of:

- a) a polypeptide comprising the amino acid sequence of SEQ ID NO:1,
- b) a polypeptide comprising a naturally occurring amino acid sequence having at least 90% sequence identity to the amino acid sequence of SEQ ID NO:1,
- c) a biologically active fragment of a polypeptide having the amino acid sequence of SEQ ID NO:1, wherein said biologically active fragment has serine ~~dehydratase~~ hydratase activity, and
- d) an immunogenic fragment of a polypeptide having the amino acid sequence of SEQ ID NO:1, wherein said immunogenic fragment generates an antibody that specifically binds to SEQ ID NO:1, the method comprising:
 - i) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding said polypeptide, and
 - ii) recovering the polypeptide so expressed.

10. (Withdrawn.) An isolated antibody which specifically binds to a polypeptide of claim 1.

11. (Previously presented.) An isolated polynucleotide selected from the group consisting of:

- a) a polynucleotide comprising the polynucleotide sequence of SEQ ID NO:2,
- b) a polynucleotide comprising a naturally occurring polynucleotide sequence at least 90% identical to the polynucleotide sequence of SEQ ID NO:2,
- c) a polynucleotide completely complementary to a polynucleotide of a),
- d) a polynucleotide completely complementary to a polynucleotide of b), and
- e) an RNA equivalent of a)-d).

12. (Canceled.)

13. (Withdrawn.) A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and
- b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

14. (Withdrawn.) A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

15. (Withdrawn.) A method of detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and

- b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

16. (Withdrawn.) A composition comprising a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

17. (Withdrawn.) A composition of claim 16, wherein the polypeptide has an amino acid sequence of SEQ ID NO:1.

18. (Withdrawn.) A method for treating a disease or condition associated with decreased expression of functional SDHH, comprising administering to a patient in need of such treatment the composition of claim 16.

Claims 19-26: (Canceled)

27. (Withdrawn.) A method of screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,
- b) detecting altered expression of the target polynucleotide, and
- c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

28. (Withdrawn.) A method of assessing toxicity of a test compound, the method comprising:

- a) treating a biological sample containing nucleic acids with the test compound,
- b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions

whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof,

- c) quantifying the amount of hybridization complex, and
- d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

Claims 29-46: (Canceled)